

The VSOP Survey: Final individual results

R. Dodson^{* a†}, S. Horiuchi^b, W. Scott^c, E. Fomalont^d, Z. Paragi^e, S. Frey^f, K. Wiik^g, H. Hirabayashi^a, P. Edwards^h, Y. Murata^a, G. Moellenbrock^d, L. Gurvits^e, and S. Tingay^b

^aISAS, JAXA, 3-1-1 Yoshinodai, Sagamihara, Japan

^bCentre for Astrophysics and Supercomputing, University of Swinburne, Melbourne, Australia

^cPhysics and Astronomy Department, University of Calgary, Canada

^dNational Radio Astronomy Observatory, Charlottesville, U.S.

^eJoint Institute for VLBI in Europe, Dwingeloo, Netherlands

^fFÖMI Satellite Geodetic Observatory, Budapest, Hungary

^gTuorla Observatory, Piikkiö, Finland

^hAustralia Telescope National Facility, CSIRO, Sydney, Australia

E-mail: rdodson@vsop.isas.jaxa.jp

In February 1997 the Japanese radio astronomy satellite HALCA was launched to provide the space-borne element for the VSOP mission. HALCA provided linear baselines three-times greater than that of ground arrays, thus providing higher resolution and higher AGN brightness temperature measurements and limits. Twenty-five percent of the scientific time of the mission was devoted to the “VSOP survey” of bright, compact, extra-galactic radio sources at 5 GHz. A complete list of 294 survey targets were selected from pre-launch surveys, 91% of which were observed during the satellite’s lifetime. The major goals of the VSOP Survey are statistical in nature: to determine the brightness temperature and approximate structure, to provide a source list for use with future space VLBI missions, and to compare radio properties with other data throughout the electromagnetic spectrum. All the data collected have now been analysed and is being prepared for the final image Survey paper. In this paper we present details of the mission, images of the sources, and some statistics of the contributions and completeness.

*The 8th European VLBI Network Symposium
September 26-29, 2006
Toruń, Poland*

^{*}Speaker.

[†]Present address: Marie Curie Fellow, OAN, Alcalá de Henares, España




Figure 1: a) A histogram of the contribution of all 28 antennae used to make the VSOP survey images, for Paper III, V and both. b) The final status of all the VSOP survey targets.

1. Introduction

The VLBI Space Observing Program (VSOP) satellite HALCA provided the space baseline for the observation of a complete sample of bright compact Active Galactic Nuclei (AGNs) at 5 GHz; the VSOP survey [2]. Of this set of 294 AGNs: 102 were presented in [4] (hereafter P-III), 140 will be presented in [1] (hereafter P-V), and 29 were not observed. The remaining 23 did not produce space fringes, where we expected to find them. These maybe the true result, however we have erred on the side of caution and consider them to be failures. Figure 1a) plots the contribution of all the antennae to the published, or to be published, experiments, and Figure 1b) plots the final status of all the survey experiments.

The images produced are compared, where possible, to both the General Observing Time (GOT), and any other available, images. We confirmed, as reported in [3], that the Survey data reproduces that of the more completely sampled images. See Figure 2 which plots all the images from P-V.

2. Conclusions

We have completed the VSOP survey imaging data reduction. The paper covering 140 sources is in preparation, and when it is published the imaging portion of the VSOP Survey Project will be completed. All the images from the final paper are shown in a montage.

Acknowledgments

The contributions of very many people have been essential to the success of the VSOP Survey project. Many more than can be included in the list of authors. We wish to recognise and thank them for their contributions.

References

- [1] Dodson, R., et al., 2007, *in preparation*, P-V
- [2] Hirabayashi H., et al., 2000, PASJ, 52, 997, P-I
- [3] Lister M. L., Piner B. G., Tingay S. J., 2000, *Astrophysical Phenomena Revealed by SVLBI*, Proceedings of the VSOP Symposium, eds: H. Hirabayashi, P. G. Edwards, & D. W. Murphy, p. 189
- [4] Scott W. K., et al., 2004, ApJS, 155, 33, P-III




Figure 2: A montage of all the images which will appear in the VSOP Survey Paper V.

The VSOP Survey: Final aggregate results

Richard Dodson^{1†}, S. Horiuchi^{2,3,4}, W. Scott⁵, E. Fomalont⁶,
 Z. Paragi⁷, S. Frey⁸, K. Wiik^{1,9}, H. Hirabayashi¹, P. Edwards^{1,10},
 Y. Murata¹, G. Moellenbrock¹¹, L. Gurvits⁷, Z. Shen¹², J. Lovell¹⁰

¹The Institute of Space and Astronautical Science, JAXA, 3-1-1 Yoshinodai, 229-8510, Japan

²Centre for Astrophysics and Supercomputing, Swinburne University, Vic. 3122, Australia

³National Astronomical Observatory, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

⁴Jet Propulsion Laboratory, 4800 Oak Grove Drive, Pasadena, CA 91109, USA

⁵Physics and Astronomy Department, University of Calgary, Calgary, Canada, T2N 1N4

⁶National Radio Astronomy Observatory, 520 Edgemont Road, Charlottesville, VA 22903, USA

⁷Joint Institute for VLBI in Europe, P.O. Box 2, 7990 AA, Dwingeloo, Netherlands

⁸FÖMI Satellite Geodetic Observatory, P.O. Box 546, H-1373, Budapest, Hungary

⁹Tuorla Observatory, Väisäläntie 20, FIN-21500 Piikkiö, Finland

¹⁰Australia Telescope National Facility, CSIRO, P. O. Box 76, Epping NSW 2122, Australia

¹¹National Radio Astronomy Observatory, P.O. Box 0, Socorro, NM 87801, USA

¹²Shanghai Astronomical Observatory, Chinese Academy of Sciences, 80 Nandan Lu, China

Abstract.

In February 1997 the Japanese radio astronomy satellite HALCA was launched to provide the space-borne element for the VSOP mission. HALCA provided linear baselines three-times greater than that of ground arrays, thus providing higher resolution and higher AGN brightness temperature measurements and limits. Twenty-five percent of the scientific time of the mission was devoted to the “VSOP survey” of bright, compact, extra-galactic radio sources at 5 GHz. A complete list of 294 survey targets were selected from pre-launch surveys, 91% of which were observed during the satellite’s lifetime. The major goals of the VSOP Survey are statistical in nature: to determine the brightness temperature and approximate structure, to provide a source list for use with future space VLBI missions, and to compare radio properties with other data throughout the electro-magnetic spectrum. All the data collected have now been analysed and is being prepared for the final image Survey paper. In this paper we present details of the mission, and some statistics of the images and brightness temperatures.

Keywords. surveys, galaxies: active

1. Introduction

The VLBI Space Observing Program (VSOP) satellite HALCA provided the space baseline for the observation of a complete sample of bright compact Active Galactic Nuclei (AGNs) at 5 GHz; the VSOP survey (Hirabayashi *et al.* 2000). Of this set of 294 AGNs: 102 were presented in Scott *et al.* (2004) (hereafter P-III), 140 will be presented in Dodson *et al.* (2007) (hereafter P-V), and 29 were not observed. The remaining 23 did not produce space fringes, where we expected to find them. Some of these may be correct, however we have erred on the side of caution and consider them to be failures.

The HALCA downlink had a bandwidth of 128 Mbps, or 32 MHz at Nyquist, two bit, sampling. A typical VSOP Survey experiment would consist of three Ground Radio Telescopes (GRTs) and a single tracking pass of HALCA; normally several hours. This leads to a nominal fringe sensitivity of 100mJy, and a image noise level of 10mJy, for the space baselines. Figure 1a) shows the number of antennae in each experiment.

† Present address: Marie Curie Fellow, OAN, Alcalá de Henares, España




Figure 1. a) A histogram of the number of antennae for experiments in P-III, V and both. b) A histogram of the upper limits to the source rest frame brightness temperatures from the imaged data in P-V, in logarithmic bins.

One of the core goals of the VSOP Survey imaging program was to measure lower limits to the brightness temperatures (T_b) directly from the data (c.f. Horiuchi *et al.* 2004). The determined T_b (from VLBI) depends only on the physical baseline length, independent of frequency, so space baselines will always provide the highest possible limits. Figure 1b) plots the lower limits to T_b for the sources in P-V, binned logarithmically. These are the source frame T_b values, unless the redshift is unknown. In which case the observer frame values are used, as a lower limit. Lower limits were estimated from the lowest brightness temperature model compatible with the data.

2. Conclusions

We have completed the VSOP survey imaging data reduction. The paper covering 140 sources is in preparation, and when it is published the imaging portion of the VSOP Survey Project will be completed. We have directly measured the source brightness temperatures, and produce a distribution of the lower limits to directly measured T_b .

Acknowledgements

RD, KJW and JEJL acknowledge support from the Japan Society for the Promotion of Science. WKS wishes to acknowledge support from the Canadian Space Agency. SH acknowledges support through an NRC/NASA-JPL Research Associateship. SF acknowledges the Bolyai Scholarship received from the Hungarian Academy of Sciences. RD wishes to thank S. Lorenzo. The NRAO is a facility of the National Science Foundation, operated under cooperative agreement by Associated Universities, Inc. The Australia Telescope Compact Array is part of the Australia Telescope which is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO.

References

- Dodson, R., *et al.*, 2007, *in preparation*, P-V
- Hirabayashi H., *et al.*, 2000, *PASJ*, 52, 997, P-I
- Horiuchi S., *et al.*, 2004, *ApJ*, 616, 110, P-IV
- Scott W. K., *et al.*, 2004, *ApJS*, 155, 33, P-III